

# New Zealand National Immunisation Schedule from 1 January 2017

	DTaP-IPV- HepB/Hib	PCV13	RV5	Hib	MMR	DTaP-IPV	Tdap	HPV	Td	Influenza
6 weeks	Infanrix <sup>®</sup> -hexa	Prevenar 13 <sup>®</sup>	RotaTeq <sup>®</sup>							
3 months	Infanrix <sup>®</sup> -hexa	Prevenar 13 <sup>®</sup>	RotaTeq <sup>®</sup>							
5 months	Infanrix <sup>®</sup> -hexa	Prevenar 13 <sup>®</sup>	RotaTeq <sup>®</sup>							
15 months		Prevenar 13 <sup>®</sup>		Act-HIB <sup>®</sup>	M-M-R <sup>®</sup> II					
4 years					M-M-R <sup>®</sup> II	Infanrix <sup>®</sup> -IPV				
11 years (school year 7)							Boostrix <sup>®</sup>			
12 years (school year 8)								Gardasil <sup>®</sup> or Gardasil <sup>®</sup> 9		
								Gardasil <sup>®</sup> or Gardasil <sup>®</sup> 9		
Every pregnancy							Boostrix <sup>®</sup>			Influvac <sup>®</sup>
45 years									ADT <sup>™</sup> Booster	
65 years									ADT <sup>™</sup> Booster	Influvac <sup>®</sup>

**Vaccine key** – **DTaP-IPV-HepB/Hib**: diphtheria, tetanus, acellular pertussis, polio, hepatitis B, Hib; **PCV13**: pneumococcal conjugate vaccine (13 valent); **RV5**: rotavirus; **Hib**: *Haemophilus influenzae* type b; **MMR**: measles, mumps, rubella; **DTaP-IPV**: diphtheria, tetanus, acellular pertussis, polio; **Tdap**: tetanus, diphtheria, acellular pertussis; **HPV**: human papillomavirus; **Td**: tetanus, diphtheria.

# Funded vaccines for special groups from 1st January 2017

Please refer to individual vaccines for detailed eligibility criteria and to the electronic *Immunisation Handbook 2014 (3rd edition)* for vaccine administration schedules

<b>Asplenia – Functional or Pre- or Post-Splenectomy Immunisation Programme</b> <ul style="list-style-type: none"> <li>Hib, influenza, meningococcal, pneumococcal, and Tdap vaccines</li> </ul>	<b>Influenza Immunisation Programme</b> <ul style="list-style-type: none"> <li>» Pregnancy,</li> <li>» Children aged 6 months to under 5 years who have been hospitalised for respiratory illness or have a history of significant respiratory illness,</li> <li>» Individuals aged 6 months to under 65 years with an eligible medical condition,</li> <li>» Individuals aged 65 years or older.</li> </ul> <ul style="list-style-type: none"> <li>Influenza vaccine</li> </ul> <b>Kidney disease</b> <ul style="list-style-type: none"> <li>Hepatitis B, Hib, influenza, pneumococcal, Tdap, and varicella vaccines</li> </ul> <b>Liver disease</b> <ul style="list-style-type: none"> <li>Hepatitis A and varicella vaccines</li> </ul> <b>Meningococcal disease case – contact with</b> <ul style="list-style-type: none"> <li>Meningococcal vaccine</li> </ul> <b>Needle stick injury – following</b> <ul style="list-style-type: none"> <li>Hepatitis B vaccine</li> </ul> <b>Non-consensual sexual intercourse – following</b> <ul style="list-style-type: none"> <li>Hepatitis B vaccine</li> </ul> <b>Pneumococcal disease – increased risk</b> <ul style="list-style-type: none"> <li>Additional pneumococcal vaccines</li> </ul> <b>Pregnancy</b> <ul style="list-style-type: none"> <li>Influenza and Tdap vaccines in every pregnancy</li> </ul> <b>Rubella – women of childbearing age who are not immune to rubella</b> <ul style="list-style-type: none"> <li>MMR vaccine</li> </ul> <b>Solid organ transplantation</b> <b>Prior to solid organ transplantation</b> <ul style="list-style-type: none"> <li>Hib, meningococcal, pneumococcal, Tdap, and varicella vaccines</li> </ul> <b>Following solid organ transplantation</b> <ul style="list-style-type: none"> <li>Hepatitis A, hepatitis B, Hib, HPV, influenza, meningococcal, pneumococcal, and Tdap vaccines</li> </ul> <b>Tuberculosis – infants and children aged under 5 years at risk of tuberculosis (TB) exposure</b> <ul style="list-style-type: none"> <li>BCG vaccine</li> </ul>
<b>Chemotherapy – following</b> <ul style="list-style-type: none"> <li>Hib, HPV, influenza, pneumococcal, Tdap, and varicella vaccines</li> </ul> <b>Also consider immunosuppression for longer than 28 days</b> <ul style="list-style-type: none"> <li>Hepatitis B and meningococcal vaccines</li> </ul>	
<b>Cochlear implant</b> <ul style="list-style-type: none"> <li>Hib, influenza, and pneumococcal vaccines</li> </ul>	
<b>Error of metabolism at risk of major metabolic decompensation</b> <ul style="list-style-type: none"> <li>Influenza and varicella vaccines</li> </ul>	
<b>Haematopoietic stem cell transplantation (HSCT) – following</b> <ul style="list-style-type: none"> <li>Hib, HPV, influenza, meningococcal, pneumococcal, Tdap, and varicella vaccines</li> </ul> <b>Also consider immunosuppression for longer than 28 days</b> <ul style="list-style-type: none"> <li>Hepatitis B vaccine</li> </ul>	
<b>Hepatitis A case – contact with</b> <ul style="list-style-type: none"> <li>Hepatitis A vaccine</li> </ul>	
<b>Hepatitis B case – contact with</b> <b>Infants born to mothers who are hepatitis B surface antigen (HBsAg) positive</b> <ul style="list-style-type: none"> <li>Hepatitis B vaccine and hepatitis B immunoglobulin (HBIG) at birth</li> </ul> <b>Household and sexual contacts of known acute hepatitis B cases or carriers</b> <ul style="list-style-type: none"> <li>Hepatitis B vaccine</li> </ul>	
<b>Hepatitis C positive individual</b> <ul style="list-style-type: none"> <li>Hepatitis B vaccine</li> </ul>	
<b>HIV positive individual</b> <ul style="list-style-type: none"> <li>Hepatitis B, HPV, influenza, meningococcal, pneumococcal, and varicella vaccines</li> </ul>	
<b>Immune deficiency/immunosuppression</b> <b>Individuals with an immune deficiency</b> <ul style="list-style-type: none"> <li>Influenza, meningococcal, and pneumococcal vaccines</li> </ul> <b>Household contacts of children or adults who will be/are immunosuppressed</b> <ul style="list-style-type: none"> <li>Varicella vaccine</li> </ul> <b>Prior to elective immunosuppression for longer than 28 days</b> <ul style="list-style-type: none"> <li>Varicella vaccine</li> </ul> <b>Following immunosuppression for longer than 28 days</b> <ul style="list-style-type: none"> <li>Hepatitis B, Hib, influenza, meningococcal, and Tdap vaccines</li> </ul>	